

UNIVERSITY OF TORONTO
STUDIES

PAPERS FROM THE PHYSICAL
LABORATORIES

NO. 74: ON THE EXTREME ULTRA-VIOLET SPECTRA OF
MAGNESIUM AND SELENIUM, by J. C. McLENNAN, J. F. L.
YOUNG, AND H. J. C. BRETON

(ABSTRACTED FROM TRANSACTIONS OF THE ROYAL SOCIETY OF CANADA, SERIES III, VOL. XIV)

THE UNIVERSITY LIBRARY: PUBLISHED BY
THE LIBRARIAN, 1920

University of Toronto Studies
COMMITTEE OF MANAGEMENT

Chairman SIR ROBERT ALEXANDER, LL.D., K.C.M.G.
President of the University

PROFESSOR J. J. ALEXANDER, Ph.D.

PROFESSOR J. J. MACKENZIE, B.A., M.B.

PROFESSOR J. P. McMURRICIE, Ph.D.

BRIG.-GEN. C. H. MITCHELL, B.A., SC., C.B., C.M.A., D.S.O.

PROFESSOR G. H. NEEDLER, Ph.D.

PROFESSOR GEORGE M. WRONG, M.A.

General Editor H. H. LANGTON, M.A.
Librarian of the University



On the Extreme Ultra-violet Spectra of Magnesium and Selenium

By PROFESSOR J. C. McLENNAN, F.R.S., MR. J. F. T. YOUNG, M.A.
and Mr. H. J. C. IRETON, M.A.

(Read May Meeting, 1919.)

INTRODUCTION

The present work is a continuation of that carried out by two of the authors with a Hilger quartz spectrograph, Type C, in the region down to 2000 A.U. The present paper deals with the spark and arc emission spectra of magnesium from 2100 A.U. to 1850 A.U. with a Hilger Type A quartz spectrograph, and the vacuum arc spectra of magnesium and selenium from 2300 A.U. to 1400 A.U. with a specially constructed fluorite spectrograph.

In the spark spectrum of magnesium, four new lines have been observed, in the arc spectrum of this metal, eleven, and in the arc spectrum of selenium, some thirty-one.

MAGNESIUM

Experiments

Three sources of radiation, the spark in air, the arc in air, and the arc in vacuo, were used in obtaining the spectra of magnesium. The spark in air was produced by the condensed discharge of a Clapp-Eastham half-kilowatt transformer, giving 10,000 volts at the secondary terminals. With this apparatus a strong thick spark was easily produced.

The arc in air was produced by using rods of magnesium metal in the carbon holders of an ordinary hand-fed arc lamp. The voltage applied was 200 volts, and the current varied from eight to twelve amperes. With such heavy currents the metal soon became hot and exposures had to be intermittent to allow for cooling of the rods.

The type of arc lamp developed for vacuum work by McLennan, Ainslie and Fuller¹ was employed with the fluorite spectrograph, magnesium metal rods being fastened to the electrodes. With good vacua it was found that a current of 4 to 5 amperes at 100 volts produced a brilliant and steady arc.

¹ Proc. Roy. Soc. Ser. A. Vol. 95. Mar. 15, 1919, p. 316.

The spectra were recorded on Schumann plates made by the Adam Hilger Co. With the quartz spectrograph no difficulty prevented the securing of clear, sharply defined spectrograms, but with the fluorite spectrograph, the fluorescence of the prism and lenses produced a heavy general fogging of the plate. Since this did not occur with other arc sources of equal intensity in the visible region, it is possible that this may be due to the strong ultra-violet emission in the magnesium arc between 2700 A.U. and 3000 A.U.

Typical spectra are reproduced in Fig. 1. The upper spectrum is that of the magnesium spark in air, and below it is the spectrum of the magnesium arc in air, both taken with the quartz spectrograph. The lower illustration is the spectrum of the magnesium arc in vacuo taken with the fluorite spectrograph. Wave length scales are attached for reference.

It may be noted that as was observed by Saunders¹ and as mentioned in the previous work, reversal was readily obtained with the arc in air at 2852 A.U. and 2026 A.U., the frequencies of which are given by $\nu = (1.5, S) - (2, P)$ and $\nu = (1.5, S) - (3, P)$, while no reversals were obtained with the arc in vacuo.

The wave lengths with the quartz spectrograph were obtained from a calibration curve constructed by using the following prominent lines in the mercury arc and the zinc, cadmium and aluminium sparks.

Mercury	Zinc	Cadmium	Aluminium
4046.78 A.U.	3076 A.U.	3260.1 A.U.	1990.57 A.U.
3650.31 "	2558.2 "	2748.68 "	1935.90 "
3341.70 "	2502.2 "	2573.15 "	1862.81 "
3131.66 "	2138.7 "	2313.88 "	1854.80 "
	2100.06 "	2288.12 "	
	2062.08 "	2265.04 "	
	2025.51 "	2194.71 "	
		2144.44 "	

The calibration curve for the fluorite spectrograph was obtained by measurement of the following prominent carbon, tin and lead vacuum arc lines.

¹ Saunders. *Astro.-Phys. Jl.* Vol. 43, No. 3. April, 1916.

Carbon	Tin	Lead
2307.5 A.U.	1756.6 A.U.	2204.4 A.U.
1930.5 "		1821.7 "
1656.9 "		1796.5 "
1562.0 "		26.5 "
61.2 "		1682.5 "
60.5 "		71.6 "
1464.5 "		1555.8 "
		1434.0 "

The wave lengths of the lines and their relative intensities are given in Table I, together with lines recorded by other observers in the region below $\lambda=2026$ A.U. In reaching these values, a great many plates were taken and the best measured with a Hilger comparator.

SERIES RELATIONS

In the present investigation the series of single lines given by the formula $\nu = (1.5, S) - (m, P)$.

m =	2	3	4	5	6	7
$\lambda =$	2853.22	2025.08	1828.1	1748.09	1707.3	1683.64

has been verified for the values $m = 3, 4, 5, 6$.

Lorenser¹ has calculated the wavelengths of the different members of the series $\nu = (1.5, S) - (m, p_2)$ to be

¹ Lorenser, *Mang. Diss.* Tübingen, 1913.

TABLE I

HANDKE		LYMAN		SAUNDERS		THE AUTHOR'S			
Spark in Air	Int.	Spark in Hydrogen	Int.	Vacuum Arc	Int.	Spark in Air	Int.	Arc in Air	Int.
..		..		2026 A.U.	10	2026 A.U.	10	2026 A.U.	10
..		..		..		2000	1	2004	1
..		..		..		1990	1	2000	1
..		..		..		1978	1	1996	1
..		..		..		1931		..	
1930.9 A.U.	6	..		..		1909	10	1973	1
1886.8	5	..		..		1887	2	..	
1864.1	4	..		..		1864	4	..	
55.9	5	..		..		1856	2	..	
39.6	3	..		..		..	2	..	
1753.0	3	1828.1 A.U.	1	1828.0 A.U.	6	..		..	
1750.7	2	1753.6	6	..		..		1828	5
50.0	1	50.9	5	..		..		1753	3
1746.7	1	..		..		..		51	1
44.1	5	..		..		..		1748	3
1741.4	5	..		..		..		..	
1736.3	2	1737.8	7	..		..		1743	1
1734.0	1	1735.0	6	..		..		1738	4
..		..		..		..		1735	1
..		..		..		..		1707	1

m =	2	3	4	5	6	7
$\lambda =$	4571.27	2090.08	1843.08	1757.1	1711.6	1621.00

Although the existence of a line at $\lambda = 2091$ A.U. was verified in the present work in both arc and spark spectra in air, no indication was obtained of the existence of members of the series with higher frequencies.

SUMMARY

1. The spectra of magnesium for spark in air and arc in air has been investigated in the region between $\lambda = 2026$ A.U. and $\lambda = 1850$ A.U. and some seven new lines recorded.

2. The spectrum of the magnesium arc in vacuo has been investigated in the region between $\lambda = 2300$ A.U. and $\lambda = 1400$ A.U. and seven new arc lines measured. Several of these agree with lines found by Saunders with the spark in hydrogen and with lines found by Handke with the spark in air.

3. The series $\nu = (1.5, S) - (m, P)$ has now been verified for values of m from two to six, but the experiments have failed to detect any further lines of the series $\nu = (1.5, S) - (m, p_2)$ beyond, possibly, the second member.

SELENIUM

These experiments were also a continuation of previous work.¹ To secure an arc in vacuo emitting selenium radiation, solid commercial carbons were bored to a depth of one and a half inches and filled with compressed selenium metal powder. These were placed in the vacuum arc lamp previously mentioned, and a steady bright arc was obtained with a current of 4 to 6 amperes at 100 volts.

In the illustration, Fig. II, the upper spectrum is that of the carbon arc in vacuo, and the lower that of selenium in the carbon arc. The following table contains the wavelengths of all the lines measured with relative intensities. As before, several of the best of a large number of plates were measured.

¹ McLennan and Young, Phil. Mag., Vol. XXXVI, p. 450, Dec. 1918.

TABLE II

Wave-length	Int.	Element	Wave-length	Int.	Element
2307.5 A.U.	8	Carbon	1619 A.U.	1	Selenium
2165	10	Selenium	1615	2	"
38	2	"	09	1	"
2073	10	"	04	3	"
51	10	"	1591	1	"
38	10	"	85	2	"
1993	2	"	78	2	"
60	8	"	61	10	Carbon
30	8	Carbon	48	1	Carbon
17	4	Selenium	33	1	Selenium
14	4	"	31	1	"
1898	6	"	30	1	"
59	8	"	03	1	"
55	8	"	1474	1	"
1761	2	"	68	1	"
52	1	"	65	3	Carbon
1691	2	"	58	1	Selenium
75	3	"	51	1	"
71	3	"	48	1	"
56	10	Carbon	46	3	"
25	1	Selenium	37	3	"
21	1	"	30	1	"

Owing to the difficulty of securing sufficient vapour density around arcs in vacuo it was found impossible to obtain any reversals in the arc spectrum so that no conclusions can be made from the point of view of absorption about the series of selenium in this region of the ultra-violet.

SUMMARY

1. Thirty new lines have been recorded in the selenium arc spectrum between $\lambda = 2300$ A.U. and $\lambda = 1400$ A.U.

Admiralty Physical Laboratory,
South Kensington.

14th April, 1919.




Figure 1




Figure 2

UNIVERSITY OF TORONTO STUDIES

PAPERS FROM THE PHYSICAL LABORATORIES

The "Papers from the Physical Laboratories", issued as a special series of University of Toronto Studies, date from the year 1900. Nos. 1-17 were published by the Physical Department in a very limited edition and are no longer in print. For the sake of a complete record the numbering of the Papers, as forming a series of University of Toronto Studies, is made continuous with the earlier series and commences with No. 18. The earlier numbers are not now available either for sale or gift.

- | | | |
|---------|---|------|
| No. 41: | The intensity of the earth's penetrating radiation at different altitudes and a secondary radiation excited by it, by Professor J. C. McLENNAN and E. N. MACALLUM | 0.25 |
| No. 42: | On the relation between the adiabatic and isothermal Young's moduli of metals, by E. F. BURTON | 0.25 |
| No. 43: | On the rotatory dispersion of quartz, by H. F. DAWES | 0.25 |
| No. 44: | On the mobilities of ions in gases at high pressures, by A. J. DEMPSEY | 0.25 |
| No. 45: | Measurements of precision on the penetrating radiation from the earth, by ARTHUR THOMSON | 0.25 |
| No. 46: | On the number of delta particles expelled concurrently with each alpha particle emitted by polonium, by W. T. KENNEDY | 0.25 |
| No. 47: | On the mobilities of ions in air at high pressures, by Professor J. C. McLENNAN and DAVID A. KEYS | 0.25 |
| No. 48: | On the absorption spectra of mercury, cadmium, zinc, and other metallic vapours, by Professor J. C. McLENNAN and EVAN EDWARDS | 0.50 |
| No. 49: | On the infra-red emission spectrum of the mercury arc, by Professor J. C. McLENNAN and RAYMOND C. DEARLE | 0.25 |
| No. 50: | On the ionisation tracks of alpha rays in hydrogen, by Professor J. C. McLENNAN and H. N. MERCER | 0.50 |
| No. 51: | On the delta rays emitted by zinc when bombarded by alpha rays, by Professor J. C. McLENNAN and C. G. FOUNDA | 0.25 |
| No. 52: | On the ultra-violet spectrum of elementary silicon by Professor J. C. McLENNAN and EVAN EDWARDS | 0.25 |
| No. 53: | On the ionisation potentials of magnesium and other metals, and on their absorption spectra, by Professor J. C. McLENNAN | 0.25 |
| No. 54: | On the Bunsen flame spectra of metallic vapours, by Professor J. C. McLENNAN and ANDREW THOMSON | 0.25 |
| No. 55: | On the ionisation of metallic vapours in flames, by Professor J. C. McLENNAN and DAVID A. KEYS | 0.25 |
| No. 56: | Emission and absorption in the infra-red spectrum of mercury, by RAYMOND C. DEARLE | 0.25 |
| No. 57: | Some experiments on residual ionization, by K. H. KINGDON | 0.25 |

No. 58:	Image-formation by Crystalline media, by H. F. DAWES.	0.25
No. 59:	Polyadic precipitation, by Miss A. W. FOSTER.	0.25
No. 60:	Regularities in the spectra of lead and tin, by R. V. ZIMMER.	0.25
No. 61:	New lines in the extreme ultra-violet of certain metals, by D. S. AXSFORD and D. S. PETER.	0.25
No. 62:	The absorption of helium by charcoal, by STEPHEN McLENNAN.	0.25
No. 63:	On the extreme-violet spectra of thallium, aluminium, lead and tin, and arsenic, by J. C. McLENNAN, J. F. T. YOUNG, and H. J. C. IRETON.	0.25
No. 64:	On the optical transparency of certain specimens of boron, by J. F. T. YOUNG and H. J. C. IRETON.	0.25
No. 65:	Use of the helium photometer for the estimation of small amounts of helium or hydrogen in air, by J. C. McLENNAN and R. T. EDWARDS.	0.25
No. 66:	The estimation of the helium content of mixtures of gases by the use of a Katharometer, by V. F. MURRAY.	0.25
No. 67:	The permeability of balloon fabrics to hydrogen and to helium, by R. T. EDWARDS and V. F. MURRAY.	0.25
No. 68:	A continuous flow apparatus for the purification of oxygen-helium mixtures, by E. EDWARDS and R. T. EDWARDS.	0.25
No. 69:	The production of helium from the natural gases of Canada, by J. C. McLENNAN.	0.25
No. 70:	Composition of the vapour and liquid phases of the system methane-nitrogen, by H. A. McTAGGART and E. EDWARDS.	0.25
No. 71:	On the extreme ultra-violet spectra of magnesium and selenium, by J. C. McLENNAN, J. F. T. YOUNG, and H. J. C. IRETON.	0.25